

TR 85-42

ESTIMATION OF THE RATE OF A
DISCRETE-TIME MULTIVARIATE POINT PROCESS

John Gubner and Prakash Narayan¹

Electrical Engineering Department

University of Maryland

College Park, Maryland 20742

Abstract

We introduce the notion of a discrete-time, multivariate point process which can arise in the modeling of an optical communication system. We wish to estimate the rate of this process at time t given the past of the process up to time $t-1$. This requires the computation of a certain conditional expectation; we perform this computation by introducing an absolutely continuous change of measure and then applying the generalized Bayes' rule..

¹This research was sponsored by the Office of Naval Research under grant no.N0001485-G-0102 and by the Minta Martin Fund for Aerospace Research from the University of Maryland at College Park.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE OCT 1985		2. REPORT TYPE		3. DATES COVERED 00-00-1985 to 00-00-1985	
4. TITLE AND SUBTITLE Estimation of the Rate of a Discrete-Time Multivariate Point Process				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Maryland,Electrical Engineering Department,College Park,MD,20742				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT see report					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 10	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

I. Introduction

Suppose that a laser beam, whose intensity is modulated by an information source, strikes a photodetector which is a part of the receiver of an optical communication system. Suppose that the beam is also influenced by external random phenomena; this occurs, for example, if the beam passes through a turbulent atmosphere [1]. The occurrence of photoelectrons at the detector can then be modeled as a doubly-stochastic, time-space Poisson process [1]. It is often necessary to compute the conditional expectation of the rate of such a process, for example, in performing a likelihood ratio test (Snyder [2], Chapter 2 and Chapter 6). In general, the computation of this conditional expectation is quite difficult, although results are available if the rate process has a Gaussian form, and the photodetector surface is very large; in addition, results for *linear* estimates are available when the photodetector surface is arbitrary, but the rate process still has a Gaussian form [3]. (see also [4, 5] for related filtering results).

Let D represent the photosensitive surface of the photodetector. It is well known that the probability is negligible that more than one photoelectron will occur in the entire region D during a time interval of length Δt , if Δt is sufficiently small. Let D_1, \dots, D_K , be a partition of D into disjoint subregions. Let Δt be "sufficiently small," and let $n_t(k)$ denote the number of photoelectrons occurring in the region D_k , during an interval $(t, t + \Delta t]$. We should expect that for each $1 \leq k \leq K$, $n_t(k)$ takes only the values 0 or 1. In addition, we can have $n_t(k) = 1$ for at most one k ; for $i \neq k$, we must have $n_t(i) = 0$. In this paper we formulate the discrete-time version of the rate-estimation problem by introducing the notion of a discrete-time, multivariate point process. By adapting the continuous-time procedures found in (Bremaud [6], Chapter IV; see also pp. 69-70, 80 of Chapter III), we solve the problem of estimating the rate of a discrete-time, multivariate point process. (For more general discrete-time procedures, see [7, 8]).

II. Problem Statement

Let $K \geq 1$ be a fixed integer. Let $(\Omega, \mathcal{F}, \mathbf{P})$ be a probability space on which all the random variables in this section are defined. We call $\{n_t(k); t \geq 1, 1 \leq k \leq K\}$ a discrete-time, multivariate point process if each $n_t(k)$ takes only the values 0 and 1 and if the events $\{n_t(k) = 1\}, 1 \leq k \leq K$, are disjoint, so that simultaneous events do not occur. Next, let \mathcal{G}_0 denote the trivial σ -field on Ω , and set

$$\mathcal{G}_t \triangleq \sigma\{n_s(k); 1 \leq s \leq t, 1 \leq k \leq K\}.$$

Now, let \mathcal{X} be a sub- σ -field of \mathcal{F} , and let

$$\mathcal{F}_t \triangleq \mathcal{G}_t \vee \mathcal{X}; \quad t \geq 0,$$

denote the smallest σ -field containing $\mathcal{G}_t \cup \mathcal{X}$. We assume that the random variables

$$\lambda_t(k) \triangleq \mathbf{E}[n_t(k) \mid \mathcal{F}_{t-1}]; \quad t \geq 1,$$

are actually \mathcal{X} -measurable (note that $\mathcal{X} = \mathcal{F}_0$) with *known* joint distributions. For example, we might take $\mathcal{X} = \sigma(X)$ for some random variable X with known distribution function $F(x)$. Then each $\lambda_t(k)$ would be some Borel function of X . Clearly, the joint distributions of the $\{\lambda_t(k); t \geq 1, 1 \leq k \leq K\}$ would be known, at least in principle. Now, observe that $\lambda_t(k) = \mathbf{P}(n_t(k) = 1 \mid \mathcal{F}_{t-1})$; since we assume that simultaneous events do not occur,

$$\sum_{k=1}^K \lambda_t(k) = \mathbf{P}\left(\bigcup_{k=1}^K \{n_t(k) = 1\} \mid \mathcal{F}_{t-1}\right).$$

This implies that $p_t \triangleq 1 - \sum_{k=1}^K \lambda_t(k)$ is also a conditional probability, and hence, nonnegative and bounded above by 1.

Our objective is to compute

$$\hat{\lambda}_t(k) \triangleq \mathbf{E}[n_t(k) \mid \mathcal{G}_{t-1}]$$

in terms of the known joint distribution of $\{\lambda_s(k); 1 \leq s \leq t, 1 \leq k \leq K\}$.

III. A Reformulation of the Problem

We shall solve our problem by reformulating the probabilistic setting above on a different probability space, $(\bar{\Omega}, \bar{F}, \bar{Q})$. On $(\bar{\Omega}, \bar{F}, \bar{Q})$ let $\{\nu_t(k); t \geq 1, 1 \leq k \leq K\}$ be a discrete-time, multivariate point process. In a manner analogous to that outlined in the previous section, we take \bar{G}_0 to be the trivial σ -field on $\bar{\Omega}$, and set

$$\bar{G}_t \triangleq \sigma\{\nu_s(k); 1 \leq s \leq t, 1 \leq k \leq K\}; \quad t \geq 1.$$

Next, let \bar{X} be a sub- σ -field of \bar{F} , and set

$$\bar{F}_t \triangleq \bar{G}_t \vee \bar{X}; \quad t \geq 0.$$

We now assume that

$$E_Q[\nu_t(k) \mid \bar{F}_{t-1}] = \mu_t(k), \quad (1)$$

where the $\{\mu_t(k)\}$ are arbitrary constants satisfying $\mu_t(k) > 0$ and

$q_t \triangleq 1 - \sum_{k=1}^K \mu_t(k) > 0$. (The symbol E_Q denotes expectation with respect to the measure

Q). As a consequence of the above assumption, under the measure Q , each $\nu_t(k)$ is independent of \bar{G}_{t-1} . In addition, the σ -field \bar{G}_t is independent of the σ -field \bar{X} (see Appendix). Next, let the random variables $\{\bar{\lambda}_t(k); t \geq 1, 1 \leq k \leq K\}$ defined on $\bar{\Omega}$ be \bar{X} -measurable and have the same joint distributions under Q as $\{\lambda_t(k); t \geq 1, 1 \leq k \leq K\}$ (defined on Ω) under P .

Given the preceding probabilistic setting on $(\bar{\Omega}, \bar{F}, \bar{Q})$, we make the following definitions. Let

$$\begin{aligned} \bar{p}_t &\triangleq 1 - \sum_{k=1}^K \bar{\lambda}_t(k); \quad t \geq 1, \\ l_t &\triangleq \sum_{k=1}^K \left(\frac{\bar{\lambda}_t(k)}{\mu_t(k)} - \frac{\bar{p}_t}{q_t} \right) \nu_t(k) + \frac{\bar{p}_t}{q_t}; \quad t \geq 1, \end{aligned} \quad (2)$$

and

$$L_t \triangleq \prod_{s=1}^t l_s : t \geq 1,$$

with $L_0 \equiv 1$. Observe that the denominators in (2) are nonzero and deterministic. Also, $0 \leq \tilde{\lambda}_t(k) \leq 1$ \mathbf{Q} -a.s., and hence, \tilde{p}_t is clearly bounded and \mathbf{Q} -integrable. (Recall that the joint distributions of the $\{\tilde{\lambda}_t(k)\}$ under \mathbf{Q} are the same as those of the $\{\lambda_t(k)\}$ under \mathbf{P}). Consequently, l_t and L_t are \mathbf{Q} -integrable. Since $\sum_{k=1}^K \tilde{\lambda}_t(k) = 1 - \tilde{p}_t$, and $\sum_{k=1}^K \mu_t(k) = 1 - q_t$, it is easy to see that $\mathbf{E}_{\mathbf{Q}}[l_t | \tilde{\mathbf{F}}_{t-1}] = 1$. Since $L_t = L_{t-1}l_t$, it is clear that

$$\mathbf{E}_{\mathbf{Q}}[L_t | \tilde{\mathbf{F}}_{t-1}] = L_{t-1}; \quad (3)$$

i.e., L_t is an $\tilde{\mathbf{F}}_t$ -martingale under \mathbf{Q} . Since, $\mathbf{E}_{\mathbf{Q}}[L_t] = \mathbf{E}_{\mathbf{Q}}[L_1] = \mathbf{E}_{\mathbf{Q}}[l_1] = 1$, we can define a new measure $\tilde{\mathbf{P}}$ on $\tilde{\mathbf{F}}_t$ by

$$\tilde{\mathbf{P}}(F) \triangleq \int_F L_t d\mathbf{Q}; \quad F \in \tilde{\mathbf{F}}_t.$$

(Technically, we should show that the family $\{L_t\}$ is uniformly integrable. However, for our purposes this is not necessary since if we wish to compute $\mathbf{E}[\nu_t(k) | \mathcal{G}_{\tau-1}]$ for some τ , we can select any finite $T \geq \tau$ and then restrict our attention to $1 \leq t \leq T$). Observe that since $L_0 = 1$, $\tilde{\mathbf{P}} = \mathbf{Q}$ on $\tilde{\mathbf{X}}$. If $\tilde{\mathbf{E}}$ denotes expectation with respect to the measure $\tilde{\mathbf{P}}$, it is not hard to verify (since simultaneous events do not occur) that

$$\begin{aligned} \tilde{\mathbf{E}}[\nu_t(k) | \tilde{\mathbf{F}}_{t-1}] &= \mathbf{E}_{\mathbf{Q}}[\nu_t(k) \frac{L_t}{L_{t-1}} | \tilde{\mathbf{F}}_{t-1}] \\ &= \mathbf{E}_{\mathbf{Q}}[\nu_t(k) l_t | \tilde{\mathbf{F}}_{t-1}] \\ &= \tilde{\lambda}_t(k), \end{aligned} \quad (4)$$

where the first equality in (4) follows from the generalized Bayes' rule. Now, since $\tilde{\mathbf{P}} = \mathbf{Q}$ on $\tilde{\mathbf{X}}$,

$$\tilde{\lambda}_t(k) = \tilde{\mathbf{E}}[\nu_t(k) | \tilde{\mathbf{F}}_{t-1}]$$

under $\tilde{\mathbf{P}}$ is probabilistically equivalent to $\lambda_t(k)$ under \mathbf{P} . In fact, we can make the following

statement. The probabilistic relationships among

$$\{ n_t(k) \}, \{ \lambda_t(k) \}, \{ F_t \}, \{ G_t \}, \text{ and } \mathbf{X}$$

under \mathbf{P} are the same as those among

$$\{ \nu_t(k) \}, \{ \bar{\lambda}_t(k) \}, \{ \bar{F}_t \}, \{ \bar{G}_t \}, \text{ and } \bar{\mathbf{X}}$$

under $\bar{\mathbf{P}}$. In the next section we shall compute $\bar{\mathbf{E}} [\nu_t(k) \mid \bar{G}_{t-1}]$ explicitly as a Borel function of $\{ \nu_s(i); 1 \leq s \leq t-1, 1 \leq i \leq K \}$. From the statement above, it follows that $\mathbf{E} [n_t(k) \mid G_{t-1}]$ will be equal to the same Borel function applied to $\{ n_s(i); 1 \leq s \leq t-1, 1 \leq i \leq K \}$.

IV. Calculations

In this section we compute $\bar{\mathbf{E}} [\nu_t(k) \mid \bar{G}_{t-1}]$. By Bayes' rule,

$$\bar{\mathbf{E}} [\nu_t(k) \mid \bar{G}_{t-1}] = \frac{\mathbf{E}_Q [\nu_t(k) L_t \mid \bar{G}_{t-1}]}{\mathbf{E}_Q [L_t \mid \bar{G}_{t-1}]}. \quad (5)$$

Next, observe that

$$\mathbf{E}_Q [L_t \mid \bar{G}_{t-1}] = \mathbf{E}_Q [\mathbf{E}_Q [L_t \mid \bar{F}_{t-1}] \mid \bar{G}_{t-1}] = \mathbf{E}_Q [L_{t-1} \mid \bar{G}_{t-1}]$$

by equation (3). In addition,

$$\begin{aligned} \mathbf{E}_Q [\nu_t(k) L_t \mid \bar{G}_{t-1}] &= \mathbf{E}_Q [\mathbf{E}_Q [\nu_t(k) L_t \mid \bar{G}_t] \mid \bar{G}_{t-1}] \\ &= \mathbf{E}_Q [\nu_t(k) \mathbf{E}_Q [L_t \mid \bar{G}_t] \mid \bar{G}_{t-1}]. \end{aligned}$$

Hence, (5) can be rewritten as

$$\bar{\mathbf{E}} [\nu_t(k) \mid \bar{G}_{t-1}] = \frac{\mathbf{E}_Q [\nu_t(k) \mathbf{E}_Q [L_t \mid \bar{G}_t] \mid \bar{G}_{t-1}]}{\mathbf{E}_Q [L_{t-1} \mid \bar{G}_{t-1}]}. \quad (6)$$

Before proceeding further, it will be convenient to introduce the following notation. For each $t \geq 1$, let

$$\nu_t \triangleq [\nu_t(1), \dots, \nu_t(K)]' ,$$

where ' denotes transpose. Then ν_t is a K -dimensional random vector. Now, let

$$\bar{\nu}_t \triangleq [\nu_1, \dots, \nu_t]'$$

Clearly, $\bar{\nu}_t$ is a tK -dimensional random vector. Next, let

$$z_t \triangleq [z_t(1), \dots, z_t(K)]' \text{ and } \bar{z}_t \triangleq [z_1, \dots, z_t]'$$

denote "dummy" variables in \mathbb{R}^K and \mathbb{R}^{tK} , respectively. Define

$$h_t(z_t) = \sum_{k=1}^K \left\{ \frac{\lambda_t(k)}{\mu_t(k)} - \frac{p_t}{q_t} \right\} z_t(k) + \frac{p_t}{q_t}, \quad (7)$$

and

$$\bar{h}_t(z_t) = \sum_{k=1}^K \left\{ \frac{\bar{\lambda}_t(k)}{\bar{\mu}_t(k)} - \frac{\bar{p}_t}{\bar{q}_t} \right\} z_t(k) + \frac{\bar{p}_t}{\bar{q}_t}.$$

Clearly, $h_t(z_t)$ is an \mathbf{X} -measurable random variable and $\bar{h}_t(z_t)$ is an $\bar{\mathbf{X}}$ -measurable random variable. In fact, $\bar{h}_t(z_t)$ under \mathbf{Q} is probabilistically equivalent to $h_t(z_t)$ under \mathbf{P} . It is then clear that the same can be said of

$$H_t(\bar{z}_t) \triangleq \prod_{s=1}^t h_s(z_s), \quad (8)$$

and

$$\bar{H}_t(\bar{z}_t) \triangleq \prod_{s=1}^t \bar{h}_s(z_s).$$

Next observe that $l_t = \bar{h}_t(\nu_t)$, and that $L_t = \bar{H}_t(\bar{\nu}_t)$. We are now ready to compute $\mathbf{E}_{\mathbf{Q}}[L_t | \bar{\mathcal{G}}_t]$ in (6) for each $t \geq 1$. (Note that $\mathbf{E}_{\mathbf{Q}}[L_0 | \bar{\mathcal{G}}_0] = \mathbf{E}_{\mathbf{Q}}[L_0] = 1$). Observe that $L_t = \bar{H}_t(\bar{\nu}_t)$ is an $\bar{\mathbf{X}}$ -measurable function of a $\bar{\mathcal{G}}_t$ -measurable random vector. Since $\bar{\mathbf{X}}$ and $\bar{\mathcal{G}}_t$ are independent under \mathbf{Q} , it follows that

$$\mathbf{E}_{\mathbf{Q}}[\bar{H}_t(\bar{\nu}_t) | \bar{\mathcal{G}}_t] = \mathbf{E}_{\mathbf{Q}}[\bar{H}_t(\bar{z}_t)] | \bar{z}_t = \bar{\nu}_t.$$

From the remarks preceding equation (8), we have

$$\mathbf{E}_Q [L_t \mid \bar{\mathcal{G}}_t] = \mathbf{E} [H_t(\bar{z}_t) \mid \bar{z}_t = \bar{v}_t].$$

We now set

$$f_t(\bar{z}_t) \triangleq \mathbf{E} [H_t(\bar{z}_t)]. \quad (9)$$

Note that $f_t(\bar{z}_t)$ is a deterministic function of \bar{z}_t . Equation (6) becomes

$$\bar{\mathbf{E}} [\nu_t(k) \mid \bar{\mathcal{G}}_{t-1}] = \frac{\mathbf{E}_Q [\nu_t(k) f_t(\bar{v}_t) \mid \bar{\mathcal{G}}_{t-1}]}{f_{t-1}(\bar{v}_{t-1})}.$$

We can write

$$\mathbf{E}_Q [\nu_t(k) f_t(\bar{v}_t) \mid \bar{\mathcal{G}}_{t-1}] = \mathbf{E}_Q [\nu_t(k) f_t(\bar{v}_{t-1}, \nu_t) \mid \bar{\mathcal{G}}_{t-1}].$$

Now, equation (1) implies that ν_t is independent of $\bar{\mathcal{G}}_{t-1}$ under \mathbf{Q} . Therefore,

$$\mathbf{E}_Q [\nu_t(k) f_t(\bar{v}_t) \mid \bar{\mathcal{G}}_{t-1}] = \mathbf{E}_Q [\nu_t(k) f_t(\bar{z}_{t-1}, \nu_t) \mid \bar{z}_{t-1} = \bar{v}_{t-1}].$$

Since simultaneous events do not occur, it is trivial to compute

$$\begin{aligned} \mathbf{E}_Q [\nu_t(k) f_t(\bar{z}_{t-1}, \nu_t)] &= f_t(\bar{z}_{t-1}, e_k) \mathbf{Q}(\nu_t(k) = 1) \\ &= \mu_t(k) f_t(\bar{z}_{t-1}, e_k), \end{aligned}$$

where e_k is the standard unit vector in \mathbf{R}^K with a 1 in the k th position and a 0 in the other $K-1$ positions. We conclude that

$$\bar{\mathbf{E}} [\nu_t(k) \mid \bar{\mathcal{G}}_{t-1}] = \frac{\mu_t(k) f_t(\bar{v}_{t-1}, e_k)}{f_{t-1}(\bar{v}_{t-1})},$$

and hence,

$$\mathbf{E} [n_t(k) \mid \mathcal{G}_{t-1}] = \frac{\mu_t(k) f_t(\bar{n}_{t-1}, e_k)}{f_{t-1}(\bar{n}_{t-1})},$$

with the obvious meaning of the symbol \bar{n}_{t-1} .

V. Summary

We have shown that if $\{ n_t(k); t \geq 1, 1 \leq k \leq K \}$ is a discrete-time, multivariate point process residing in the probabilistic setting outlined in Section II, then

$$\mathbf{E} [n_t(k) \mid \mathcal{G}_{t-1}] = \frac{\mu_t(k) f_t(\bar{n}_{t-1}, e_k)}{f_{t-1}(\bar{n}_{t-1})},$$

where $f_t(\bar{z}_t)$ is given by equations (7), (8), and (9).

The $\mu_t(k)$ introduced in Section III were arbitrary; if we set $\mu_t(k) = \frac{1}{K+1}$ for all t and all k in the preceding equations, we find that

$$\mathbf{E} [n_t(k) \mid \mathcal{G}_{t-1}] = \frac{a_t(\bar{n}_{t-1}, e_k)}{a_{t-1}(\bar{n}_{t-1})},$$

where

$$a_t(\bar{z}_t) \triangleq \mathbf{E} \left[\prod_{s=1}^t \left(\sum_{k=1}^K (\lambda_s(k) - p_s) n_s(k) + p_s \right) \right].$$

Appendix

For $1 \leq s \leq t$, let z_s denote either the zero vector in \mathbb{R}^K or any standard unit vector in \mathbb{R}^K . In this appendix we prove that if $E \in \bar{\mathbf{X}}$, then (using the notation of Sections III and IV)

$$\mathbf{Q}(\nu_t = z_t, \dots, \nu_1 = z_1, E) = \left(\prod_{s=1}^t \mathbf{Q}(\nu_s = z_s) \right) \mathbf{Q}(E).$$

Since $\bar{\mathcal{G}}_t = \sigma\{\nu_1, \dots, \nu_t\}$, this will prove that $\bar{\mathcal{G}}_t$ and $\bar{\mathbf{X}}$ are independent under \mathbf{Q} .

Proof. Using the definition of conditional probability,

$$\mathbf{Q}(\nu_t = z_t, \dots, \nu_1 = z_1, E) = \int \mathbf{Q}(\nu_t = z_t \mid \bar{\mathbf{F}}_{t-1}) d\mathbf{Q}. \quad (10)$$

$$\{ \nu_{t-1} = z_{t-1}, \dots, \nu_1 = z_1, E \}$$

Since simultaneous events do not occur, if $z_t = e_k$, $Q(\nu_t = z_t \mid \bar{F}_{t-1}) = \mu_t(k)$. If $z_t = 0$,

$$Q(\nu_t = z_t \mid \bar{F}_{t-1}) = 1 - \sum_{k=1}^K \mu_t(k). \text{ Since } Q(\nu_t = z_t \mid \bar{F}_{t-1}) \text{ is deterministic,}$$

$$Q(\nu_t = z_t \mid \bar{F}_{t-1}) = Q(\nu_t = z_t).$$

Hence (10) becomes

$$Q(\nu_t = z_t, \dots, \nu_1 = z_1, E) = Q(\nu_t = z_t)Q(\nu_{t-1} = z_{t-1}, \dots, \nu_1 = z_1, E).$$

The remainder of the proof by induction is clear.

QED

Acknowledgement

It is a pleasure to thank Professor A. M. Makowski for several illuminating discussions on the applications of an absolutely continuous change of measure to filtering problems.

References

- [1] D. L. Snyder, "Applications of Stochastic Calculus for Point Process Models Arising in Optical Communication," pp. 789-804 in *Communication Systems and Random Process Theory*, ed. J. K. Skwirzynski, Sijthoff and Noordhoff, Alphen aan der Rijn, The Netherlands (1978).
- [2] D. L. Snyder, *Random Point Processes*, John Wiley & Sons, New York (1975).
- [3] J. Gubner and P. Narayan, "Estimation of the Rate of a Doubly-Stochastic Time-Space Poisson Process," *Systems & Control Letters*, Submitted, (1985).
- [4] D. L. Snyder and P. M. Fishman, "How to Track a Swarm of Fireflies by Observing Their Flashes," *IEEE Transactions on Information Theory*, vol. IT-21, pp. 692-695 (Nov. 1975).
- [5] I. B. Rhodes and D. L. Snyder, "Estimation and Control Performance for Space-Time Point-Process Observations," *IEEE Transactions on Automatic Control*, vol. AC-22, No. 3, pp. 338-346 (June 1977).
- [6] P. Bremaud, *Point Process and Queues, Martingale Dynamics*, Springer-Verlag, New York (1981).
- [7] P. M. Bremaud and J. H. van Schuppen, *Discrete Time Stochastic Systems: Part I: Stochastic Calculus and Representations*, Report No. 7603a, Department of Systems Science and Mathematics, Washington University, St. Louis (1976).
- [8] P. M. Bremaud and J. H. van Schuppen, *Discrete Time Processes: Part II: Estimation Theory*, Report No. 7603b, Department of Systems Science and Mathematics, Washington University, St. Louis (1976).